

## Rochester Pain Solution

### Discharge Instructions: Trigger Point

Your doctor has given you joint injection(s). This procedure is done to relieve pain, by injecting medication directly into the joint. The local anesthetic medication used will provide a temporary numbing effect. A steroid medication may be used to decrease inflammation, and will work slowly over the next one to two weeks. These effects are temporary and will diminish over time.

#### Activity:

1. After the injection you may notice increased tenderness in the area of your usual pain. For injection site tenderness, apply ice packs to the injection site for no longer than 20 minutes at a time, repeating the ice pack treatment 4 to 6 times as needed. Increase activity as tolerated.

Diet: You may resume your usual diet.

#### Medication:

1. Resume all medications as prescribed

**If you develop ANY of the following, please call the doctor you have seen for your procedure or go to the emergency room:**

- Redness, swelling, or drainage from the injection sites
- Fever or chills
- Prolonged dizziness
- Shortness of breath

**If you cannot reach your physician and it is an emergency, please call 911 or go to the nearest emergency room.**

Monday to Friday from 8:00 AM to 5:00 PM, call (585)491-6130. After 5:00PM, on weekends or Holidays, call (585)520-1812.

Follow up: You may call the office at (585)491-6130 if a follow up appointment is needed.

**If you have received intravenous sedation, we suggest you do not sign any important documents or make any important decisions for 24 hours.**

I have read and understand the above instructions and have received a copy.

Patient Signature: \_\_\_\_\_ Witness Signature: \_\_\_\_\_

Accompanying Adult: \_\_\_\_\_ Date: \_\_\_\_\_